Das tripelnegative Mammakarzinom

- eine therapeutische Herausforderung -

Priv.-Doz. Dr. med. Cornelia Liedtke
Klinik für Gynäkologie und Geburtshilfe
- Direktor: Prof. Dr. med. Achim Rody -

Universitätsklinikum Schleswig-Holstein / Campus Lübeck

Überblick

Was ist das tripelnegative Mammakarzinom?

Wie ist der klinische Verlauf?

Welche Therapien können eingesetzt werden?

Was sind die neuen Therapieansätze?

Überblick

Was ist das tripelnegative Mammakarzinom?

Wie ist der klinische Verlauf?

Welche Therapien können eingesetzt werden?

Was sind die neuen Therapieansätze?

Definition des TNBC

„Triple-negative breast cancer: disease entity

or

of clinical relevance!

~~title of convenience?”~~

Carey L, Winer E, Viale G, Cameron D, Gianni L

Nat Rev Clin Oncol 2010

Zusammenhang zwischen TNBC und BLBC

Zusammenhang zwischen TNBC und TNBC

Zusammenhang zwischen TNBC, BLBC und BRCA1

Mutationswahrscheinlichkeit beim TNBC

Patient category	% in cohort (n = 403)	% of BRCA1 carriers in each category/95 % exact binomial CI
Total	100 (403)	16.1 % (65/403)/12.7–20.1 %
“Premenopausal” (<50 years)	51.6 (208)	26.9 % (56/208)/21.0–33.5 %
Early onset (<40 years)	26.3 (106)	35.9 % (38/106)/26.8–45.7 %
Family history	26.1 (105)	47.6 % (50/105)/37.8–57.6 %
No family history, onset >51 years	43.7 (176)	2.3 % (4/176)/0.6–5.7 %
No family history, onset <50 years	30.3 (122)	9 % (11/122)/4.5–15.6 %

Fostira et al., Breast Cancer Res Treat 2012

% in BRCA1			
a. Frequency of BRCA mutations in all TNBCs^a			
	Neoadjuvant (n = 152)	Familial (n = 134)	Adjuvant (n = 91)
Mutation analysis			
BRCA1 (%)	27 (24)	0 (0)	9
BRCA2 (%)	2 (2)	0 (0)	4
BRCA1 UV (%)	2 (2)	16 (22)	0
BRCA2 UV (%)	1 (1)	7 (9)	0
Wild type (%)	80 (71)	51 (69)	0
ND	40	60	78
b. Frequency of BRCAness in all TNBCs without BRCA1/2 mutations^b			
	Neoadjuvant (n = 123)	Familial (n = 134)	Adjuvant (n = 78)
aCGH			
BRCA1 like (%)	70 (66)	92 (69)	53 (68)
Non-BRCA1 like (%)	35 (34)	42 (31)	25 (32)
ND	18	0	0
BRCA1 promoter methylation			
Methylated (%)	31 (29)	47 (37)	21 (27)
Unmethylated (%)	75 (71)	81 (63)	56 (73)
ND	17	6	1
Abbreviations: aCGH=array Comparative Genomic Hybridisation; ND=not determined; TNBC=triple-negative breast cancer; UV=unclassified variant.			
^a Patients with unknown characteristics (ND) were not included in P-value calculations; therefore percentages for these groups were not depicted in the table.			
^b For Adjuvant series, only samples with a positive mutation test result were known; therefore, we could not calculate percentages.			

Lips et al., BJC 2013

Überblick

Was ist das tripelnegative Mammakarzinom?

Wie ist der klinische Verlauf?

Welche Therapien können eingesetzt werden?

Was sind die neuen Therapieansätze?

Prognose des TNBC

Liedtke et al., J Clin Oncol 2008

Prognose des TNBC

**Aggressivere
Biologie**

**Fehlen
therapeutischer
Substanzen**

Liedtke et al., J Clin Oncol 2008

Zusammenhang zwischen molekularen Subtypen und Alter

TNBC

Liedtke et al., in prep.

Lokalisation der Metastasen beim TNBC

(Genomisches) Grading und Ansprechen auf neoadj. Chemotherapie

Liedtke et al., J Clin Oncol 2009

Überblick

Was ist das tripelnegative Mammakarzinom?

Wie ist der klinische Verlauf?

Welche Therapien können eingesetzt werden?

Was sind die neuen Therapieansätze?

Systemische Therapie bei verschiedenen molekularen Subtypen

Harbeck [...] Liedtke. Cancer Treatment Reviews 2010

Ansprechen gegenüber neoadjuvanter Chemotherapie

	n	pCR*-Rate	
		TNBC	andere
Liedtke et al.	1118	22%	11%
Bidard et al.	293	17%	4%
Fernandez-Morales et al.	100	27%	11%
Carey et al.	107	27%	11%
Kearn et al.	145	17%	3%
Rouzier et al.	82	45%**	18%**

* Pathologische Komplettremission

** basal/nicht-basal im Genexpressionsprofiling

In-vitro-Chemotherapiesensitivität bei Mammakarzinomsubtypen

Table 4 – Association between immunohistochemical breast cancer subclasses and in vitro response to chemotherapy (univariate analysis).

		Left (n = 67) 'basal-like'	Middle (n = 70) 'HER2-like'	Right (n = 279) 'luminal'
Chemosensitivity response parameters	ET	40.3	25.8	28.6
	ED	47.8	43.0	44.0
	EC	4.5	8.2	7.0

The values represent percentages of patients classified as chemosensitive or not with regard to the tested in vitro chemotherapy regimen.

Liedtke et al., Eur J Cancer 2012

Prognose in Abhängigkeit vom Chemotherapieansprechen

Anforderungen an die Forschung zum TNBC

Identifikation der molekularen Mechanismen für Chemotherapieresistenz

Identifikation von Patientinnen mit/ohne Therapieansprechen

Optimierung der Chemotherapie (-regime)

Entwicklung neuer zielgerichteter Therapien

Liedtke et al., J Clin Oncol 2008

Anforderungen an die Forschung zum TNBC

Identifikation der molekularen Mechanismen für Chemotherapieresistenz

Identifikation von Patientinnen mit/ohne Therapieansprechen

Optimierung der Chemotherapie (-regime)

Entwicklung neuer zielgerichteter Therapien

Liedtke et al., J Clin Oncol 2008

Die GeparSixto-Studie: Einsatz von Carboplatin

Von Minckwitz et al., ASCO 2013

Schwere unerwünschte Nebenwirkungen (SAEs) in der GeparSixto

	PM(+B) PMCb(+HL) (N=157)	PMCb(+B) (N=158)	PM(+HL) (N=136)
N with at least 1 SAE (%)	62 (39.5)	69 (43.7)	55 (40.4)
Infections	15	18	21
pneumonia	6	1	11
Blood	7	26	3
anemia	1	11	1
febrile neutropenia	6	6	1
Gastrointestinal	10	11	10
diarrhea	3	0	3

Von Minckwitz et al., ASCO 2013

Carboplatin neoadjuvant (CALBG 40603 / ALLIANCE)

San Antonio Breast Cancer Symposium – Cancer Therapy and Research Center at UT Health Science Center December 10-14, 2013

CALBG 40603: Schema – Randomized Phase II

2 X 2 Randomization

- Paclitaxel 80 mg/m² wkly x 12 ddAC x 4
- Paclitaxel 80 mg/m² wkly x 12 ddAC x 4 + Bevacizumab 10 mg/kg q2wks x 9
- Paclitaxel 80 mg/m² wkly x 12 ddAC x 4 + Carboplatin AUC 6 q3wks x 4
- Paclitaxel 80 mg/m² wkly x 12 ddAC x 4 + Carboplatin AUC 6 q3wks x 4 + Bevacizumab 10 mg/kg q2wks x 9

Surgery^{§*}
XRT^{*}
No Adjuvant Systemic Treatment Planned^{*}

[§]Research biopsies if residual tumor
^{*}MD discretion

Research biopsies-frozen and fixed

ALLIANCE
This presentation is the intellectual property of William Sikov, MD. Contact at wsikov@lifespan.org for permission to reprint or distribute.

pCR Breast/Axilla (ypT0/is N0) + / - Carboplatin

Group	pCR Breast and Axilla	N
No Carboplatin	41% (35-48%)	N=212
Carboplatin	54% (48-61%)	N=221

Odds ratio: 1.71
p = 0.0029

ALLIANCE
This presentation is the intellectual property of William Sikov, MD. Contact at wsikov@lifespan.org for permission to reprint or distribute.

CALBG 40603: Eligibility

- Clinical Stage II-III (except T4d – inflammatory BC)
- ER + PR ≤10%, HER(-) (IHC 0-1+ or FISH <2.0)
- No contraindication to treatment with bevacizumab
- Baseline evaluation including breast MRI (or US)
- FNA or biopsy of suspicious axillary LN recommended
- Pre-treatment SLN sampling permitted in clinical N0

Sikov et al., SABCS 2013

AGO-Empfehlungen zur PST (bei TNBC)

Neoadjuvante systemische Chemotherapie Empfohlene Regime und Schedules		Oxford / AGO LoE / GR		
> Adjuvante Standardregime mit einer Dauer von mindestens 18 Wochen		1a	A	++
> AC oder EC → D q3w oder P q1w		2b	A	++
> DAC		2b	B	++
> AP → CMF		1b	A	+
> Taxan gefolgt von Anthrazyklin		2b	B	+
> Dosisdichte Protokolle (z. B. E -P-CMF, E-P-C)		1b	B	+*
> Capecitabin in Kombination mit Anthrazyklin und Taxan		1b	B	+/-
> Platinsalze beim TNBC unabhängig von einer BRCA1-Mutation		2b	B	+*

* Studienteilnahme empfohlen

AGO 2014

Anforderungen an die Forschung zum TNBC

Identifikation der molekularen Mechanismen für Chemotherapieresistenz

Identifikation von Patientinnen mit/ohne Therapieansprechen

Optimierung der Chemotherapie (-regime)

Entwicklung neuer zielgerichteter Therapien

Liedtke et al., J Clin Oncol 2008

Effizienz von first-line-Regimen beim mTNBC

O'Shaughnessy et al., SABCS 2010; Finn et al., J Clin Oncol 2009; Baselga et al., Ann Oncol 2010; O'Shaughnessy et al. NEJM 2011; O'Shaughnessy et al., BCRT 2007; Schneeweiß et al., EMCC 2011; Aogi et al., BCRT 2011

pCR-Raten durch Bevacizumab (Gepar Quinto)

Von Minckwitz et al., NEJM 2012

Bevacizumab neoadjuvant (CALBG 40603 / ALLIANCE)

San Antonio Breast Cancer Symposium – Cancer Therapy and Research Center at UT Health Science Center December 10-14, 2013

CALBG 40603: Schema – Randomized Phase II

2 X 2
Randomization

Research biopsies-frozen and fixed

ALLIANCE
This presentation is the intellectual property of William Sikov, MD. Contact at wsikov@lifespan.org for permission to reprint or distribute.

pCR Breast/Axilla (ypT0/is N0) +/- Bevacizumab

Group	pCR Breast and Axilla (%)	95% CI	N
No Bevacizumab	44%	38-51%	218
Bevacizumab	52%	45-58%	215

Odds ratio: 1.36
p = 0.0570

ALLIANCE
This presentation is the intellectual property of William Sikov, MD. Contact at wsikov@lifespan.org for permission to reprint or distribute.

CALBG 40603: Eligibility

- Clinical Stage II-III (except T4d – inflammatory BC)
- ER + PR ≤10%, HER(-) (IHC 0-1+ or FISH <2.0)
- No contraindication to treatment with bevacizumab
- Baseline evaluation including breast MRI (or US)
- FNA or biopsy of suspicious axillary LN recommended
- Pre-treatment SLN sampling permitted in clinical N0

Sikov et al., SABCS 2013

Rezidiv-freies Überleben durch Bevacizumab beim TNBC

Überblick

Was ist das tripelnegative Mammakarzinom?

Wie ist der klinische Verlauf?

Welche Therapien können eingesetzt werden?

Was sind die neuen Therapieansätze?

Wirksamkeit von Iniparib (Phase II und III)

O'Shaughnessy et al., NEJM 2011

O'Shaughnessy et al., Proc ASCO 2011

Molekulare Heterogenität des TNBC

PIK3CA →

EGFR →

Shah et al., Nature 2012

Definition verschiedener Subtypen des TNBC

TNBC subtype	Cell line	Subtype correlation (P value)	Histology	Mutations ^A	Intrinsic subtype ^B	Basal subtype ^C
Basal-like						
BL1	HCC2157	0.66 (0.00)	DC	<i>BRCA1; STAT4; UTX</i>	Basal	Basal A
	HCC1599	0.62 (0.00)	DC	<i>BRCA2; TP53; CTNND1; TOP2B; CAMK1G</i>	Basal	Basal A
	HCC1937	0.28 (0.00)	DC	<i>BRCA1; TP53; MAPK13; MDC1</i>	HER2	Basal A
	HCC1143	0.26 (0.00)	IDC	<i>TP53</i>	Basal	Basal A
	HCC3153	0.24 (0.00)		<i>BRCA1</i>	Basal	Basal A
BL2	MDA-MB-468	0.19 (0.06)	DC	<i>PTEN; RB1; SMAD4; TP53</i>	Basal	Basal A
	HCC38	0.19 (0.01)	DC	<i>CDKN2A; TP53</i>	Unclassified	Basal B
	SUM149PT	0.30 (0.00)	INF	<i>BRCA1</i>	Unclassified	Basal B
	CAL-851	0.25 (0.00)	IGA	<i>RB1; TP53</i>	Basal	Basal A
	HCC70	0.24 (0.04)	DC	<i>PTEN; TP53</i>	Basal	Basal A
IM	HCC1806	0.22 (0.00)	ASCC	<i>CDKN2A; TP53; UTX</i>	Unclassified	Basal A
	HDQ-P1	0.18 (0.17)	IDC	<i>TP53</i>	Unclassified	
	HCC1187	0.22 (0.00)	DC	<i>TP53; CTNNA1; DDX18; HUWE1; NFKBIA</i>	Basal	Basal A
	DU4475	0.17 (0.00)	DC	<i>APC; BRAF; MAP2K4; RB1</i>	Unclassified	
Mesenchymal-like						
M	BT-549	0.21 (0.00)	IDC	<i>PTEN; RB1; TP53</i>	Unclassified	Basal B
	CAL-51	0.17 (0.00)	AC	<i>PIK3CA</i>	Unclassified	
	CAL-120	0.09 (0.00)	AC	<i>TP53</i>	Luminal B	
MSL	HS578T	0.29 (0.00)	CS	<i>CDKN2A; HRAS; TP53</i>	Unclassified	Basal B
	MDA-MB-157	0.25 (0.00)	MBC	<i>NF1; TP53</i>	Unclassified	Basal B
	SUM159PT	0.14 (0.00)	ANC	<i>PIK3CA; TP53 HRAS</i>	Unclassified	Basal B
	MDA-MB-436	0.13 (0.00)	IDC	<i>BRCA1; TP53</i>	Unclassified	Basal B
	MDA-MB-231	0.12 (0.00)	IDC	<i>BRAF; CDKN2A; KRAS; NF2; TP53; PDGFRA</i>	Unclassified	Basal B
LAR						
LAR	MDA-MB-453	0.53 (0.00)	AC	<i>PIK3CA; CDH1; PTEN</i>	Luminal A	Luminal
	SUM185PE	0.39 (0.00)	DC	<i>PIK3CA</i>	Luminal A	Luminal
	HCC2185	0.34 (0.00)		<i>PIK3CA</i>	Luminal A	Luminal
	CAL-148	0.32 (0.00)	AC	<i>PIK3CA; RB1; TP53; PTEN</i>	Luminal A	
	MFM-223	0.31 (0.00)	AC	<i>PIK3CA; TP53</i>	Luminal A/B	
Unclassified	HCC1395		DC	<i>ATR; BRCA2; CDKN2A; PTEN; FGFR1; PDGFRB; TP53</i>	Basal	
	BT20		IDC	<i>CDKN2A; PIK3CA; TP53</i>	HER2	Basal A
	SW527				Luminal B	

Lehmann et al., JCI 2011

Differentielles Ansprechen auf Chemotherapie

The relationship between RCB index and the 7 TNBC subtypes

	BL1	BL2	M	IM	MSL	LAR	UNS
pCR	11	0	8	8	3	2	5
RCB-I	1	1	2	3	2	4	1
RCB-II	6	6	4	8	6	8	3
RCB-III	3	1	9	5	2	4	6

Symmans WF et al. J Clin Oncol 2007

PRESENTED AT: ASCO Annual Meeting 2013

Masuda et al., ASCO 2013

Das Immunsystem – eine Achillesferse?!

Definition von TILs

San Antonio Breast Cancer Symposium
- Cancer Therapy and Research Center at UT Health Science Center - December 10-14, 2013

Predefined parameters for TIL evaluation

intratumoral TILs = direct contact to tumor cells

stromal TILs = between the tumor cells

LPBC = Lymphocyte-predominant breast cancer
„more lymphocytes than tumor cells“ (≥60% TILs)

Denkert C, Loibl S, et al. J Clin Oncol. 2010, Issa-Nummer et al. PLOS

This presentation is the intellectual property of the author/presenter. Contact them at carsten.denkert@charite.de for permission to reprint and/or distribute.

San Antonio Breast Cancer Symposium
- Cancer Therapy and Research Center at UT Health Science Center - December 10-14, 2013

20-28% of patients hat a Lymphocyte-predominant breast cancer (LPBC) phenotype

LPBC: „more lymphocytes than tumor cells“ (≥60% stromal or intratumoral TILs)

Group	n	LPBC Prevalence
all tumors	580	24%
triple-negative tumors	314	28%
HER2+ tumors	266	20%

the intellectual property of the author/presenter. Contact them at ert@charite.de for permission to reprint and/or distribute.

Denkert et al., SABCS

Prognostische Wertigkeit der B-Zell / IL-8 -Signatur beim TNBC

Überblick

Was ist das tripelnegative Mammakarzinom?

Wie ist der klinische Verlauf?

Welche Therapien können eingesetzt werden?

Gibt es unterschiedliche Subgruppen?

Herzlichen Dank für Ihre Aufmerksamkeit ...

Wissen schafft Gesundheit

Klinik für Gynäkologie und Geburtshilfe

Safe the Date ...

**Lübeck
28.06.2014**